arXiv:dg-ga/9712021v1 30 Dec 1997

On the Spinor Representation of Surfaces in Euclidean 3-Space. *

Thomas Friedrich, Berlin

February 7, 2008

1 Introduction

The Weierstraß formula describes a conformal minimal immersion of a Riemann surface M^2 into the 3-dimensional Euclidean space $\mathbb{R}^3$. It expresses the immersion in terms of a holomorphic function g and a holomorphic 1-form μ as the integral

$$f = \mathrm{Re}\left(\int (1 - g^2, i(1 + g^2), 2g)\mu \right) : M^2 \to \mathbb{R}^3.$$

On the other hand let us consider the spinor bundle S over M^2. This 2-dimensional vector bundle splits into

$$S = S^+ \oplus S^- = \Lambda^0 \oplus \Lambda^{1,0}.$$

Therefore the pair (g, μ) can be considered as a spinor field φ on the Riemann surface. The Cauchy-Riemann equation for g and μ is equivalent to the homogeneous Dirac equation

$$D(\varphi) = 0.$$

The choice of the Riemannian metric in the fixed conformal class of M^2 is not essential since the kernel of the Dirac operator is a conformal invariant.

A similar description for *an arbitrary* surface $M^2 \hookrightarrow \mathbb{R}^3$ is possible and has been pointed out probably for the first time by Eisenhardt (1909). This representation of any surface in $\mathbb{R}^3$ by a spinor field φ on M^2 satisfying the inhomogeneous Dirac equation

$$D(\varphi) = H\varphi \qquad (*)$$

involving the mean curvature H of the surface has been used again in some recent papers (see [KS], [R], [Tai1], [Tai2], [Tai3]). However, the mentioned authors describe the relationship between surfaces in $\mathbb{R}^3$ and solutions of the equation $(*)$ in local terms in order to get explicit formulas. The aim of the present paper is to clarify the mentioned representation of surfaces in $\mathbb{R}^3$ by solutions of the equation $D(\varphi) =$

---

*Supported by the SFB 288 of the DFG.

$H\varphi$ in a *geometrically invariant way.* It turns out that the main idea leading to the description of a surface by a spinor field φ is simple: Consider an immersion $M^2 \hookrightarrow \mathbb{R}^3$ and fix a parallel spinor Φ on $\mathbb{R}^3$. Then the restriction $\varphi = \Phi_{|M^2}$ of Φ to the surface is (with respect to the inner geometry of M^2) a non-trivial spinor field on M^2 and defines a spinor φ^* of constant length which is a solution of the inhomogeneous Dirac equation

$$D(\varphi^*) = H\varphi^*.$$

Conversely, given a solution φ of the equation $(*)$ with constant length there exists a symmetric endomorphism $E : T(M^2) \to T(M^2)$ such that the spinor field satisfies a "twistor type equation"

$$\nabla_X \varphi = E(X) \cdot \varphi.$$

The resulting integrability conditions for the endomorphism E are exactly the Gauß and Codazzi equations. As a consequence, the solution φ of the Dirac equation

$$D(\varphi) = H\varphi \quad , \quad |\varphi| \equiv const > 0$$

yields an isometric immersion of M^2 into $\mathbb{R}^3$. In a similar way one obtains the description of conformal immersions using the well-known formula for the transformation of the Dirac operator under a conformal change of the metric (see [BFGK]).

2 The Dirac Operator of a Surface immersed into a Riemannian 3-Manifold.

Let Y^3 be a 3-dimensional oriented Riemannian manifold with a fixed spin structure and denote by M^2 an oriented surface isometrically immersed into Y^3. Because the normal bundle of M^2 is trivial, the spin structure of Y^3 induces a spin structure on the Riemannian surface M^2. The spinor bundle S of the 3-manifold Y^3 yields by restriction the spinor bundle of the surface M^2. Over M^2 this bundle decomposes into

$$S = S^+ \oplus S^-$$

where the subbundles $S^\pm$ are defined by (see [F])

$$S^\pm = \{\varphi \in S : i \cdot e_1 \cdot e_2 \cdot \varphi = \pm\varphi\}$$

Here $\{e_1, e_2\}$ denotes an oriented orthonormal frame in $T(M^2)$ and $X \cdot \varphi$ means the Clifford multiplication of a spinor $\varphi \in S$ by a vector $X \in T(M^2)$. Since in the 3-dimensional Clifford algebra the relation

$$e_1 \cdot e_2 = e_3$$

holds, we can replace the Clifford product $e_1 \cdot e_2$ by the normal vector $\vec{N}$ of $M^2 \hookrightarrow Y^3$:

$$S^\pm = \{\varphi \in S : i \cdot \vec{N} \cdot \varphi = \pm\varphi\}.$$

Consider a spinor field Φ defined on the 3-manifold Y^3. Its restriction $\varphi = \Phi_{|M^2}$ is a spinor field defined on M^2 and decomposes therefore into $\varphi = \varphi^+ + \varphi^-$ with

$$\varphi^+ = \frac{1}{2}(\varphi + i\vec{N} \cdot \varphi) \quad , \quad \varphi^- = \frac{1}{2}(\varphi - i\vec{N} \cdot \varphi).$$

We denote by ∇^{Y^3} and ∇^{M^2} the covariant derivatives in the spinor bundles on Y^3 and M^2 respectively. For any vector $X \in T(M^2)$ we have the well-known formula (see [BFGK])

$$\nabla^{Y^3}_X(\Phi) = \nabla^{M^2}_X(\varphi) - \frac{1}{2}(\nabla_X \vec{N}) \cdot \vec{N} \cdot \varphi.$$

The vector $(\nabla_X \vec{N})$ coincides with the second fundamental form II: $T(M^2) \to T(M^2)$ of the submanifold $M^2 \hookrightarrow Y^3$. Since II is symmetric the Clifford product $e_1 \cdot \mathrm{II}(e_1) + e_2 \cdot \mathrm{II}(e_2)$ is a scalar and equals $(-2H)$ where H denotes the mean curvature of the surface M^2. The Dirac operator D of M^2 defined by the formula

$$D(\varphi) = e_1 \cdot \nabla^{M^2}_{e_1} \varphi + e_2 \cdot \nabla^{M^2}_{e_2} \varphi$$

can now be expressed by the covariant derivative ∇^{Y^3} and the mean curvature vector:

$$e_1 \cdot \nabla^{Y^3}_{e_1}(\Phi) + e_2 \cdot \nabla^{Y^3}_{e_2}(\Phi) = D(\varphi) + H \cdot \vec{N} \cdot \varphi.$$

Suppose that the spinor field Φ on Y^3 is a real Killing spinor, i.e. there exists a number $\lambda \in \mathbb{R}^1$ such that for any tangent vector $\vec{T} \in T(Y^3)$ the derivative of Φ in the direction of $\vec{T}$ is given by the Clifford multiplication:

$$\nabla^{Y^3}_{\vec{T}}(\Phi) = \lambda \cdot \vec{T} \cdot \Phi.$$

For the restriction $\varphi = \Phi_{|M^2}$ we obtain immediately the equation

$$D(\varphi) = -2\lambda\varphi - H \cdot \vec{N} \cdot \varphi.$$

Using the decomposition $\varphi = \varphi^+ + \varphi^-$ the last equation is equivalent to the pair of equations

$$D(\varphi^+) = (-2\lambda - iH)\varphi^- \quad , \quad D(\varphi^-) = (-2\lambda + iH)\varphi^+.$$

We discuss two special cases.

Proposition 1: *Let M^2 be a minimal surface in Y^3. Then the restriction $\varphi = \Phi_{|M^2}$ of any real Killing spinor Φ on Y^3 is an eigenspinor of constant length on the surface M^2:*

$$D(\varphi) = -2\lambda\varphi.$$

On the other hand, suppose that Φ is a parallel spinor ($\lambda = 0$) on Y^3. Then we obtain

$$D(\varphi^+) = -iH\varphi^- \quad , \quad D(\varphi^-) = iH\varphi^+.$$

If we introduce the spinor field $\varphi^* = \varphi^+ - i\varphi^-$, a simple calculation shows

$$D(\varphi^*) = H\varphi^*.$$

The spinor field φ^* is given by

$$\varphi^* = \varphi^+ - i\varphi^- = \frac{1}{2}(\varphi + i \cdot \vec{N} \cdot \varphi) - \frac{i}{2}(\varphi - i \cdot \vec{N} \cdot \varphi) = \frac{1}{2}(1-i)\varphi + \frac{1}{2}(-1+i) \cdot \vec{N} \cdot \varphi.$$

Moreover, the length of φ^* is constant. This construction yields the

Proposition 2: *Let Φ be a parallel spinor field defined on the 3-manifold Y^3 and denote by $\varphi = \Phi_{|M^2}$ its restriction to M^2. Define the spinor field φ^* on M^2 by the formula*

$$\varphi^* = \frac{1}{2}(1-i)\varphi + \frac{1}{2}(-1+i) \cdot \vec{N} \cdot \varphi.$$

Then φ^* is a spinor field of constant length on M^2 satisfying the Dirac equation

$$D(\varphi^*) = H\varphi^*$$

where H denotes the mean curvature.

Remark 1: The map $\Phi \longmapsto \varphi^*$ associating to any parallel spinor Φ on Y^3 a solution of the equation $D(\varphi^*) = H\varphi^*$ is injective.

Remark 2: We can apply the above mentioned formulas not only for Killing spinors. Indeed, for any spinor field Φ we have

$$D_{Y^3}(\Phi) = D(\varphi) + H \cdot \vec{N} \cdot \varphi + \vec{N} \cdot (\nabla^{Y^3}_{\vec{N}} \Phi)$$

where D_{Y^3} is the Dirac operator of the 3-manifold Y^3. Suppose there exists a function $\kappa : M^2 \to \mathbb{C}$ such that the normal derivative $(\nabla^{Y^3}_{\vec{N}} \Phi)$ of the spinor field Φ is described by κ:

$$\left(\nabla^{Y^3}_{\vec{N}} \Phi\right) = \kappa\Phi.$$

Then we obtain

$$D_{Y^3}(\Phi) = D(\varphi) + (H + \kappa)\vec{N} \cdot \varphi.$$

This formula (in arbitrary dimension) has been used for the calculation of the spectrum of the Dirac operator on hypersurfaces of the Euclidean space (see [Bä], [Tr1], [Tr2]).

3 Solutions of the Dirac Equation with Potential on Riemannian Surfaces.

Let (M^2, g) be an oriented, 2-dimensional Riemannian manifold with spin structure. $H : M^2 \to \mathbb{R}^1$ denotes a given smooth, real-valued function defined on the surface. In this part we study spinor fields φ on M^2 that are solutions of the differential equation

$$D(\varphi) = H\varphi.$$

If we decompose the spinor field into $\varphi = \varphi^+ + \varphi^-$ according to the splitting $S = S^+ \oplus S^-$ of the spinor bundle the equation we want to study is equivalent to the system

$$D(\varphi^+) = H\varphi^- \quad , \quad D(\varphi^-) = H\varphi^+.$$

To any solution φ of this equation we associate two forms $F_\pm$ defined for pairs $X, Y \in T(M^2)$ of tangent vectors:

$$F_+(X,Y) = \mathrm{Re}(\nabla_X \varphi^+, Y \cdot \varphi^-) \quad , \quad F_-(X,Y) = \mathrm{Re}(\nabla_X \varphi^-, Y \cdot \varphi^+).$$

Proposition 3:

a.) $F_\pm$ are symmetric bilinear forms on $T(M^2)$.

b.) The trace of $F_\pm$ is given by $Tr(F_\pm) = -H|\varphi^\mp|^2$.

Proof: The symmetry of $F_\pm$ is a consequence of the Dirac equation as well as the assumption that H is a real-valued function. Indeed, we have

$$\begin{aligned}
\mathrm{Re}(\nabla_{e_1}\varphi^+, e_2\varphi^-) &= \mathrm{Re}(e_1 \cdot \nabla_{e_1}\varphi^+, e_1 \cdot e_2 \cdot \varphi^-) = \mathrm{Re}(H\varphi^- - e_2 \cdot \nabla_{e_2}\varphi^+, e_1 \cdot e_2 \cdot \varphi^-) = \\
&= H \cdot \mathrm{Re}(\varphi^-, e_1 \cdot e_2 \cdot \varphi^-) + \mathrm{Re}(\nabla_{e_2}\varphi^+, e_2 \cdot e_1 \cdot e_2 \cdot \varphi^-) = \\
&= 0 + \mathrm{Re}(\nabla_{e_2}\varphi^+, e_1 \cdot \varphi^-).
\end{aligned}$$

Moreover, we calculate the trace of $F_\pm$:

$$Tr(F_\pm) = \mathrm{Re}(\nabla_{e_1}\varphi^\pm, e_1 \cdot \varphi^\mp) + \mathrm{Re}(\nabla_{e_2}\varphi^\pm, e_2 \cdot \varphi^\mp) = -\mathrm{Re}(D(\varphi^\pm), \varphi^\mp) = -H|\varphi^\mp|^2.$$

■

We study now special solutions of the equation $D(\varphi) = H\varphi$, i.e. solutions with *constant length* $|\varphi| \equiv$ *const* $\neq 0$. It may happen that the components $\varphi^\pm$ have a non-empty zero set.

Proposition 4: *Suppose that the spinor field φ defined on the Riemannian surface M^2 is a solution of the equation*

$$D(\varphi) = H\varphi \qquad \textit{with} \ \ |\varphi| \equiv const \neq 0.$$

Then the forms $F_\pm$ are related by the equation

$$|\varphi^+|^2 F_+ = |\varphi^-|^2 F_- .$$

Proof: In case one of the spinors φ^+ or φ^- vanishes at a fixed point $m_0 \in M^2$ the relation between F_+ and F_- is trivial. Otherwise there exists a neighbourhood V of the point $m_0 \in M^2$ such that both spinors φ^+ and φ^- are not zero at any point $m \in V$. The spinors

$$\frac{e_1 \cdot \varphi^-}{|\varphi^-|} \quad , \quad \frac{e_2 \cdot \varphi^-}{|\varphi^-|}$$

are an orthonormal base in S^+ with respect to the Euclidean scalar product $\mathrm{Re}(\cdot,\cdot)$. Therefore we obtain (on V)

$$\begin{aligned} \nabla_X \varphi^+ &= \mathrm{Re}\left(\nabla_X \varphi^+, \frac{e_1 \cdot \varphi^-}{|\varphi^-|}\right) \frac{e_1 \cdot \varphi^-}{|\varphi^-|} + \mathrm{Re}\left(\nabla_X \varphi^+, \frac{e_2 \cdot \varphi^-}{|\varphi^-|}\right) \frac{e_2 \cdot \varphi^-}{|\varphi^-|} = \\ &= \frac{1}{|\varphi^-|^2}\{F_+(X,e_1)e_1 + F_+(X,e_2)e_2\} \cdot \varphi^- . \end{aligned}$$

A similar calculation yields the formula

$$\nabla_X \varphi^- = \frac{1}{|\varphi^+|^2}\{F_-(X,e_1)e_1 + F_-(X,e_2)e_2\} \cdot \varphi^+ .$$

We multiply the equations by φ^+ and φ^- respectively and sum up. Then we obtain

$$\frac{1}{2}\nabla_X(|\varphi^+|^2 + |\varphi^-|^2) = \mathrm{Re}(A(X)\varphi^-, \varphi^+)$$

where the endomorphism $A : T(M^2) \to T(M^2)$ is defined by

$$A(X) = \left\{ \frac{F_+(X,e_1)}{|\varphi^-|^2} - \frac{F_-(X,e_1)}{|\varphi^+|^2} \right\} e_1 + \left\{ \frac{F_+(X,e_2)}{|\varphi^-|^2} - \frac{F_-(X,e_2)}{|\varphi^+|^2} \right\} e_2 .$$

Since $F_\pm$ are symmetric tensors, the endomorphism A is symmetric too. Moreover, the trace of A vanishes:

$$TrA = \frac{1}{|\varphi^-|^2} Tr(F_+) - \frac{1}{|\varphi^+|^2} Tr(F_-) = -H + H = 0.$$

The length of the spinor field φ is constant. This implies

$$\mathrm{Re}(A(X) \cdot \varphi^-, \varphi^+) = 0.$$

At any point $m \in V$ of the set V the spinors φ^+, φ^- are non-trivial. Then the rank of the endomorphisms $A : T(M^2) \to T(M^2)$ is not greater than one. All in all, A is symmetric, $Tr(A) = 0$ and $rg(A) \leq 1$, i.e. $A \equiv 0$.

■

We now consider the sum

$$F = F_+ + F_- .$$

At points with $\varphi^+ \neq 0$ (or $\varphi^- \neq 0$) we have

$$\frac{F}{|\varphi|^2} = \frac{F_+ + F_-}{|\varphi^+|^2 + |\varphi^-|^2} = \frac{\left(\frac{|\varphi^-|^2}{|\varphi^+|^2} + 1\right) F_-}{|\varphi^+|^2 + |\varphi^-|^2} = \frac{F_-}{|\varphi^+|^2}$$

as well as

$$\frac{F}{|\varphi|^2} = \frac{F_+ + F_-}{|\varphi^+|^2 + |\varphi^-|^2} = \frac{\left(1 + \frac{|\varphi^+|^2}{|\varphi^-|^2}\right) F_+}{|\varphi^+|^2 + |\varphi^-|^2} = \frac{F_+}{|\varphi^-|^2}.$$

The endomorphism $E : T(M^2) \to T(M^2)$, $E = \frac{F}{|\varphi|^2}$, is defined at all points of M^2 and the formulas derived in the proof of proposition 4 in fact prove the following

Proposition 5: *Let φ be a solution of the differential equation $D(\varphi) = H\varphi$ on a Riemannian surface (M^2, g) with a real-valued function $H : M^2 \to \mathbb{R}^1$. Suppose that the length $|\varphi| \equiv const \neq 0$ of the spinor field φ is constant. Then*

$$E(X) = \frac{1}{|\varphi|^2}\mathrm{Re}(\nabla_X \varphi, Y \cdot \varphi)$$

is a symmetric endomorphism $E : T(M^2) \to T(M^2)$ such that

a.) $\nabla_X \varphi^+ = E(X) \cdot \varphi^- \quad , \quad \nabla_X \varphi^- = E(X) \cdot \varphi^+$

b.) $Tr(E) = -H$.

■

For a given triple (M^2, g, E) of a Riemannian surface and symmetric endomorphism the existence of a non-trivial solution φ of the equation

$$\nabla_X \varphi = E(X) \cdot \varphi$$

implies certain integrability conditions. It turns out that in this way we obtain precisely the well-known Gauß and Codazzi equations of the classical theory of surfaces in Euclidean 3-space.

Proposition 6: *Let (M^2, g) be a 2-dimensional Riemannian surface with a fixed spin structure and suppose that $E : T(M^2) \to T(M^2)$ is a symmetric endomorphism. If there exists a non-trivial solution of the equation*

$$\nabla_X \varphi = E(X) \cdot \varphi \quad , \quad X \in T(M^2)$$

then

a.) (Codazzi equation): $\nabla_X(E(Y)) - \nabla_Y(E(X)) - E([X,Y]) = 0.$

b.) (Gauß equation): $\det(E) = \frac{1}{4}G$, *where G is the Gaussian curvature of (M^2, g).*

Proof: We prove the two equations in a way similar to the derivation of the integrability conditions for the Riemannian metric in case the space admits a Killing spinor (see [BFGK]). We differentiate the equation

$$\nabla_X \varphi = E(X) \cdot \varphi$$

and then we calculate the curvature tensor R^S of the spinor bundle S:

$$\begin{aligned} R^S(X,Y)\varphi &= \nabla_X \nabla_Y \varphi - \nabla_Y \nabla_X \varphi - \nabla_{[X,Y]}\varphi = \\ &= \{\nabla_X(E(Y)) - \nabla_Y(E(X)) - E([X,Y]) + E(Y)E(X) - E(X)E(Y)\} \cdot \varphi. \end{aligned}$$

On the other side, the curvature tensor $R^S : S \to S$ is given by the formula

$$R^S(e_1, e_2) = \frac{1}{2} R_{1212}\ e_1 \cdot e_2.$$

Denote by $A(X,Y)$ the differential of E:

$$A(X,Y) = \nabla_X(E(Y)) - \nabla_Y(E(X)) - E([X,Y]).$$

A simple algebraic calculation in the spin representation then leads to the equations

$$-A(X,Y)\varphi^- = \left(2\det(E) + \frac{R_{1212}}{2}\right) i\varphi^+$$

$$A(X,Y)\varphi^+ = \left(2\det(E) + \frac{R_{1212}}{2}\right) i\varphi^-.$$

We multiply the first equation once by the vector $A(X;Y)$:

$$||A(X,Y)||^2 \varphi^- = -\left(2\det(E) + \frac{R_{1212}}{2}\right)^2 \varphi^-$$

and then we conclude $A(X,Y) \equiv 0$ (Codazzi equation) as well as $\det(E) = -\frac{1}{4}R_{1212} = \frac{1}{4}G$ (Gauß equation).

■

For a given triple (M^2, g, E) consisting of a Riemannian spin surface (M^2, g) and of a symmetric endomorphism E we will denote by $\mathcal{K}(M^2, g, E)$ the space of all spinor fields φ satisfying the equation $\nabla_X \varphi = E(X) \cdot \varphi$. It is invariant under the quaternionic structure $\alpha : S \to S$, i.e. $\mathcal{K}(M^2, g, E)$ is a quaternionic vector space (see §4). Denote by $(-H)$ the trace of E,

$$Tr(E) = -H.$$

Then we have

$$\mathcal{K}(M^2, g, E) \subset \ker(D - H).$$

In this part of the paper we prove that any spinor field $\varphi \in \ker(D - H)$ of constant length belongs to one of the subspaces $\mathcal{K}(M^2, g, E)$ for a suitable symmetric endomorphism E, $Tr(E) = -H$.

Finally, we consider the lengths

$$L_+ = ||\varphi^+||^2 \quad , \quad L_- = ||\varphi^-||^2$$

of a non-trivial solution $\varphi \in \mathcal{K}(M^2, g, E)$. Using the integrability condition $\det(E) = \frac{G}{4}$ (i.e. $||E||^2 = H^2 - \frac{G}{2}$) as well as the well-known formula $D^2 = \Delta + \frac{G}{2}$ for the square D^2 of the Dirac operator we can derive formulas for $\Delta(L_\pm)$:

$$\begin{aligned}
\Delta(L_\pm) &= 2(\Delta(\varphi^\pm), \varphi^\pm) - 2\langle\nabla(\varphi^\pm), \nabla(\varphi^\pm)\rangle \\
&= 2(D^2(\varphi^\pm), \varphi^\pm) - 2\left(\frac{G}{2}\right) \bullet ||\varphi^\pm||^2 - 2||E||^2||\varphi^\mp||^2 \\
&= 2\left(H^2 - \frac{G}{2}\right)(L_\pm - L_\mp) + 2\mathrm{Re}(grad\,(H) \cdot \varphi^\mp, \varphi^\pm)
\end{aligned}$$

In particular, if $H \equiv const$ is constant, the difference $u = L_+ - L_-$ satisfies the differential equation

$$\Delta(u) = 4\left(H^2 - \frac{G}{2}\right)u.$$

4 The Period Form of a Spinor with $\nabla_X\varphi = E(X) \cdot \varphi$.

We consider a spinor field φ on a Riemannian surface (M^2, g) such that

$$\nabla_X\varphi = E(X) \cdot \varphi$$

for a fixed symmetric endomorphism E. The spinor bundle S carries a quaternionic structure $\alpha : S \to S$ commuting with Clifford multiplication and interchanging the decomposition $S = S^+ \oplus S^-$ (see [F]). For any spinor field $\varphi = \varphi^+ + \varphi^-$ we define three 1-forms by

$$\xi^\varphi(X) = 2(X \cdot \varphi^+, \varphi^-)$$

$$\xi_+^\varphi(X) = (X \cdot \varphi^+, \alpha(\varphi^+)) \quad , \quad \xi_-^\varphi(X) = (X \cdot \varphi^-, \alpha(\varphi^-)).$$

ξ^φ and ξ_+^φ are $\Lambda^{1,0}$-forms, ξ_-^φ is a $\Lambda^{0,1}$-form. Indeed, $e_1 \cdot e_2$ acts on S^+ (on S^-) by multiplication by $(-i)$ (by i). Now we obtain

$$(\star\xi^\varphi)(e_1) = -\xi^\varphi(e_2) = 2(-e_2 \cdot \varphi^+, \varphi^-) = (-ie_2 \cdot e_1 \cdot e_2 \cdot \varphi^+, \varphi^-) = -i\xi^\varphi(e_1),$$

i.e. $\star\xi^\varphi = -i\xi^\varphi$ holds. A similar calculation gives $\star\xi_+^\varphi = -i\xi_+^\varphi$ and $\star\xi_-^\varphi = i\xi_-^\varphi$. We split the 1-form ξ^φ into its real and imaginary part:

$$\xi^\varphi = w^\varphi + i\mu^\varphi.$$

Moreover, we introduce the 1-form Ω^φ

$$\Omega^\varphi = \xi_+^\varphi - \xi_-^\varphi.$$

Then we have

Proposition 7: *Let (M^2, g) be a Riemannian spin surface and $E : T(M^2) \to T(M^2)$ a symmetric endomorphism of trace $-H$. Suppose the spinor field φ is a solution of the equation $\nabla_X\varphi = E(X) \cdot \varphi$. Then* a.) $dw^{\varphi} = 0$.

b.) $d\mu^{\varphi} = 2H\{|\varphi^-|^2 - |\varphi^+|^2\}dM^2$.

c.) $d\Omega^{\varphi} = 0$.

Proof: We calculate dw^{φ}:

$$\begin{aligned}
\frac{1}{2}dw^{\varphi}(X,Y) &= X(\mathrm{Re}(Y\cdot\varphi^+,\varphi^-)) - Y(\mathrm{Re}(X\cdot\varphi^+,\varphi^-)) - \mathrm{Re}([X,Y]\cdot\varphi^+,\varphi^-) = \\
&= \{g(X,E(Y)) - g(Y,E(X))\}|\varphi^-|^2 + \{g(X,E(Y)) - g(Y,E(X))\}|\varphi^+|^2.
\end{aligned}$$

Since E is symmetric, we obtain $dw^{\varphi} = 0$. A similar calculation shows the formula for $d\mu^{\varphi}$. For the proof of $d\Omega^{\varphi} = 0$ we first remark that the quaternionic structure $\alpha : S \to S$ and the hermitian product $(\cdot,\cdot)$ on S are related by

$$(\varphi_1, \alpha(\varphi_2)) = -\overline{(\alpha(\varphi_1), \varphi_2)}.$$

Using this formula we can transform $d\xi_-^{\varphi}$ in the following way:

$$\begin{aligned}
d\xi_-^{\varphi}(X,Y) &= (Y\cdot E(X)\cdot\varphi^+, \alpha(\varphi^-)) + (Y\cdot\varphi^-, \alpha(E(X)\cdot\varphi^+)) \\
&\quad -(X\cdot E(Y)\cdot\varphi^+, \alpha(\varphi^-)) - (X\cdot\varphi^-, \alpha(E(Y)\cdot\varphi^+)) \\
&= -\overline{(\alpha(Y\cdot E(X)\cdot\varphi^+), \varphi^-)} - (E(X)\cdot Y\cdot\varphi^-, \alpha(\varphi^+)) \\
&\quad +\overline{(\alpha(X\cdot E(Y)\cdot\varphi^+), \varphi^-)} + (E(Y)\cdot X\cdot\varphi^-, \alpha(\varphi^+)) \\
&= -(E(X)\cdot Y\cdot\varphi^-, \alpha(\varphi^+)) - (E(X)\cdot Y\cdot\varphi^-, \alpha(\varphi^+)) \\
&\quad +(E(Y)\cdot X\cdot\varphi^-, \alpha(\varphi^+)) + (E(Y)\cdot X\cdot\varphi^-, \alpha(\varphi^+)).
\end{aligned}$$

On the other hand we calculate $d\xi_+^{\varphi}$:

$$\begin{aligned}
d\xi_+^{\varphi}(X,Y) &= (Y\cdot E(X)\cdot\varphi^-, \alpha(\varphi^+)) + (Y\cdot\varphi^+, \alpha(E(X)\cdot\varphi^-)) \\
&\quad -(X\cdot E(Y)\cdot\varphi^-, \alpha(\varphi^+)) - (X\cdot\varphi^+, \alpha(E(Y)\cdot\varphi^-)) = \\
&= (Y\cdot E(X)\cdot\varphi^-, \alpha(\varphi^+)) - (E(X)\cdot Y\cdot\varphi^+, \alpha(\varphi^-)) \\
&\quad -(X\cdot E(Y)\cdot\varphi^-, \alpha(\varphi^+)) + (E(Y)\cdot X\cdot\varphi^+, \alpha(\varphi^-)).
\end{aligned}$$

Finally we obtain

$$\begin{aligned}
d(\xi_-^{\varphi} - \xi_+^{\varphi})(X,Y) &= -(\{E(X)\cdot Y + Y\cdot E(X)\}\varphi^-, \alpha(\varphi^+)) \\
&\quad +(\{E(Y)\cdot X + X\cdot E(Y)\}\varphi^-, \alpha(\varphi^+)) - \\
&= 2\{g(E(X),Y) - g(E(Y),X)\}(\varphi^-, \alpha(\varphi^+))
\end{aligned}$$

and $d(\xi_-^{\varphi} - \xi_+^{\varphi}) = 0$ follows again by the symmetry of E.

■

Let us consider the case that (M^2, g) is isometrically immersed into the Euclidean space $\mathbb{R}^3$, Φ is a parallel spinor on $\mathbb{R}^3$ and the spinor field φ^* on M^2 defined by the formula

$$\varphi^* = \frac{1}{2}\left(\Phi_{|M^2} + i\cdot\vec{N}\cdot\Phi_{|M^2}\right) + \frac{i}{2}\left(i\cdot\vec{N}\cdot\Phi_{|M^2} - \Phi_{|M^2}\right)$$

(see §2). In this case the forms w^{φ^*} and Ω^{φ^*} are given by the expressions

$$w^{\varphi^*}(X) = -\mathrm{Im}(X\cdot\Phi,\Phi) \quad , \quad \Omega^{\varphi^*}(X) = (X\cdot\Phi,\alpha(\Phi)),$$

and are exact 1-forms. Indeed, we defined functions $f:\mathbb{R}^3\to\mathbb{R}^1$ and $g:\mathbb{R}^3\to\mathbb{C}$ by

$$f(m) = -\mathrm{Im}\langle m\cdot\Phi,\Phi\rangle \quad , \quad g(m) = \langle m\cdot\Phi,\alpha(\Phi)\rangle$$

and then we have $df = w^{\varphi^*}, dg = \Omega^{\varphi^*}$. We remark that f and g describe in fact the isometric immersion $M^2 \hookrightarrow \mathbb{R}^3$ we started with. The 3-dimensional spinor $\Phi\in\Delta_3$ defines a real 3-dimensional subspace $\Delta_3(\Phi)$ by

$$\Delta_3(\Phi) = \{\Psi\in\Delta_3 : \quad \mathrm{Re}(\Psi,\Phi) = 0\}.$$

The map $\Psi \to (-\mathrm{Im}(\Psi,\Phi),(\Psi,\alpha(\Phi)))$ is an isometry between $\Delta_3(\Phi)$ and $\mathbb{R}^1\oplus\mathbb{C} = \mathbb{R}^3$. Clearly, the immersion $M^2\hookrightarrow\mathbb{R}^3$ is given by

$$M^2 \ni m \longmapsto m\cdot\Phi \in \Delta_3(\Phi),$$

i.e. by the functions $f_{|M^2}$ and $g_{|M^2}$. With respect to $d(f_{|M^2}) = w^{\varphi^*}$ and $d(g_{|M^2}) = \Omega^{\varphi^*}$ we obtain a formula for the isometric immersion $M^2\hookrightarrow\mathbb{R}^3$:

$$\oint (w^{\varphi^*},\Omega^{\varphi^*}) : M^2 \to \mathbb{R}^3.$$

(Weierstraß representation of the surface.)

In general, we call a solution φ of the differential equation $\nabla_X\varphi = E(X)\cdot\varphi$ *exact* iff the corresponding forms w^φ,Ω^φ are exact 1-forms. Using the definition

$$Hess\,(h)(X,Y) = \frac{1}{2}\{g(\nabla_X(grad\,(h)),Y) + g(X,\nabla_Y(grad\,(h)))\}$$

of the Hessian of a smooth function h defined on a Riemannian manifold we obtain the

Proposition 8: *Let $\varphi\in\mathcal{K}(M^2,g,E)$ be an exact solution of the differential equations $\nabla_X\varphi = E(X)\cdot\varphi$ with $df = w^\varphi$, $dg = \Omega^\varphi$. Then*

a.) $Hess\,(f) = 2\left(|\varphi^+|^2 - |\varphi^-|^2\right)E.$

b.) $|grad\,f|^2 = 4|\varphi^+|^2|\varphi^-|^2.$

c.) $Hess\,(g) = -4(\varphi^-,\alpha(\varphi^+))E.$

d.) $|grad\,(g)|^2 = (|\varphi^+|^2 - |\varphi^-|^2)^2.$

■

In particular, the determinant of the Hessian of the function f is given by

$$\det(Hess\,(f)) = 4(|\varphi^+|^2 - |\varphi^-|^2)^2 \det(E) = (|\varphi^+|^2 - |\varphi^-|^2)^2 G.$$

Here we used proposition 6, i.e. $\det(E) = \frac{1}{4}G$.

Corollary: *Let M^2 be a compact Riemannian spin-manifold and suppose that $\varphi \in \mathcal{K}(M^2, g, E)$ is an exact, non-trivial solution. Then the spinors φ^+ or φ^- vanish at least at out one point. Moreover, there exists $m_0 \in M^2$ such that $G(m_0) \geq 0$.*

Proof: At a maximum point $m_0 \in M^2$ of f we have

$$grad(f)(m_0) = 0 \quad , \quad \det(Hess\,(f)(m_0)) \geq 0.$$

■

Recall that for any 2-dimensional Riemannian manifold (M^2, g) and any function $h : M^2 \to \mathbb{R}^1$ the 2-form

$$\{2\det(Hess\,(h)) - |grad(h)|^2 G\} dM^2 = d\mu^1$$

is exact (see [S], page 47). Using this formula in case of an exact solution $\varphi \in \mathcal{K}(M^2, g, E)$ we obtain

$$\int\limits_{M^2} (|\varphi^+|^2 - |\varphi^-|^2)^2 G = 2 \int\limits_{M^2} |\varphi^+|^2 |\varphi^-|^2 G.$$

Corollary: *Let M^2 be a compact Riemannian spin manifold and suppose that $\varphi \in \mathcal{K}(M^2, g, E)$ is an exact solution. Then*

$$\int\limits_{M^2} (|\varphi|^4 - 6|\varphi^+|^2 |\varphi^-|^2) G = 0.$$

■

We again discuss the last formula in case of an isometrically immersed surface $M^2 \hookrightarrow \mathbb{R}^3$ and a given parallel spinor Φ on $\mathbb{R}^3$. We apply the integral formula to the spinor $\varphi^* = \varphi^*_+ + \varphi^*_-$ where

$$\varphi^*_+ = \frac{1}{2}\left(\Phi + i\vec{N} \cdot \Phi\right) \quad , \quad \varphi^*_- = \frac{1}{2}\left(-\Phi + i \cdot \vec{N} \cdot \Phi\right).$$

In this case we have

$$|\varphi^*_+|^2 = \frac{1}{2}|\Phi|^2 + \frac{1}{2}\langle i\vec{N} \cdot \Phi, \Phi\rangle \quad , \quad |\varphi^*_-|^2 = \frac{1}{2}|\Phi|^2 - \frac{1}{2}\langle i\vec{N} \cdot \Phi, \Phi\rangle$$

and ($|\Phi| \equiv 1$) therefore

$$1 - 6|\varphi^*_+|^2 |\varphi^*_-|^2 = -\frac{1}{2} + \frac{3}{2}\langle i \cdot \vec{N} \cdot \Phi, \Phi\rangle^2.$$

Consequently the integral formula yields

$$\int\limits_{M^2} G = 3 \int\limits_{M^2} \langle i\vec{N}\Phi, \Phi\rangle^2 G.$$

The spinors $i\Phi$ as well as $\vec{N} \cdot \Phi$ belong to $V(\Phi) \subset \Delta_3$, the space of the immersion $M^2 \hookrightarrow \mathbb{R}^3 = V(\Phi)$. The last formula means therefore

$$\int\limits_{M^2} G = 3 \int\limits_{M^2} \langle \vec{N}, \mathfrak{a}_3\rangle^2 G$$

for the unit vector $\mathfrak{a}_3 = i\Phi \in V(\Phi) = \mathbb{R}^3$.

5 The Spin Formulation of the Theory of Surfaces in $\mathbb{R}^3$.

An oriented, immersed surface $M^2 \hookrightarrow \mathbb{R}^3$ inherits from $\mathbb{R}^3$ an inner metric g, a spin structure and a solution φ of the Dirac equation

$$D(\varphi) = H\varphi$$

of constant length $|\varphi| \equiv 1$ where H denotes the mean curvature of the surface. The spinor field φ on M^2 is the restriction of a parallel spinor field Φ of the Euclidean space $\mathbb{R}^3$. The period forms w^φ and Ω^φ are exact and the immersion $M^2 \hookrightarrow \mathbb{R}^3$ is given by integration of the $\mathbb{R}^1 \oplus \mathbb{C} = \mathbb{R}^3$ valued form $(w^\varphi, \Omega^\varphi)$. At least locally the converse is true: Given an oriented, 2-dimensional Riemannian manifold (M^2, g) with a fixed spin structure and a solution of constant length of the Dirac equation $D(\varphi) = H\varphi$ for some smooth function $H : M^2 \to \mathbb{R}^1$, there exists a symmetric endomorphism $E : T(M^2) \to T(M^2)$ such that $\varphi \in \mathcal{K}(M^2, g, E)$. Moreover, $2E$ is the second fundamental form of an isometric immersion $(M^2, g) \to \mathbb{R}^3$. We formulate this description of the theory of surfaces in $\mathbb{R}^3$ in the following

Theorem 1: *Let (M^2, g) be an oriented, 2-dimensional Riemannian manifold and $H : M^2 \to \mathbb{R}^1$ a smooth function. Then there is a correspondence between the following datas:*

1. *an isometric immersion $(\tilde{M}^2, g) \to \mathbb{R}^3$ of the universal covering $\tilde{M}^2$ into the Euclidean space $\mathbb{R}^3$ with mean curvature H.*
2. *a solution φ with constant length $|\varphi| \equiv 1$ of the Dirac equation $D(\varphi) = H \cdot \varphi$.*
3. *a pair (φ, E) consisting of a symmetric endomorphism E such that $Tr(E) = -H$ and a spinor field φ satisfying the equation $\nabla_X \varphi = E(X) \cdot \varphi$.*

■

We apply now the well-known formulas for the change of the Dirac operator under a conformal change of the metric. Suppose that $\tilde{g} = \sigma g$ are two conformally equivalent metrics on M^2 where $\sigma : M^2 \to (0, \infty)$ is a positive function. Denote by D and $\tilde{D}$ the Dirac operator corresponding to the metric g and $\tilde{g}$ respectively. Then

$$\tilde{D}(\tilde{\varphi}) = \sigma^{-3/4} D\widetilde{(\sigma^{1/4}\varphi)}$$

holds (see [BFGK]). Let us consider a solution φ of the Dirac equation

$$D(\varphi) = \lambda\varphi$$

on (M^2, g) and suppose that φ never vanishes. We introduce the Riemannian metric $\tilde{g} = |\varphi|^4 g$ as well as the spinor field $\varphi^* = \frac{\varphi}{|\varphi|}$. Then we obtain

$$\tilde{D}(\varphi^*) = \frac{\lambda}{|\varphi|^2}\varphi^* \quad , \quad |\varphi^*| \equiv 1,$$

and thus an isometric immersion $(\tilde{M}^2, |\varphi|^4 g) \to \mathbb{R}^3$ with mean curvature $H = \frac{\lambda}{|\varphi|^2}$.

Theorem 2: *Let (M^2, g) be an oriented, 2-dimensional Riemannian manifold. Any spinor field φ without zeros that is a solution of the equation*

$$D(\varphi) = \lambda\varphi$$

defines an isometric immersion $(\tilde{M}^2, |\varphi|^4 g) \hookrightarrow \mathbb{R}^3$ with mean curvature $H = \frac{\lambda}{|\varphi|^2}$.

■

Remark 1: Consider the case that $M^2 \hookrightarrow S^3$ is a minimal surface in S^3. Let Φ be a real Killing spinor on S^3, i.e.

$$\nabla_{\vec{T}}(\Phi) = \frac{1}{2}\vec{T} \cdot \Phi.$$

The restriction $\varphi = \Phi_{|M^2}$ is an eigenspinor of the Dirac operator on M^2 with constant length (proposition 1). Therefore φ defines an isometric immersion of $(\tilde{M}^2, g) \hookrightarrow \mathbb{R}^3$ with mean curvature $H \equiv -1$. This transformation associates to any minimal surface $M^2 \hookrightarrow S^3$ a surface of constant mean curvature $H \equiv -1$ in $\mathbb{R}^3$, a well-known construction (see [La]).

Remark 2: Using the described correspondence between isometric immersions of surfaces into $\mathbb{R}^3$ and solutions of the Dirac equation $D(\varphi) = H \cdot \varphi$ one can immediately remark that several statements of the elementary theory of surfaces are equivalent to several statements concerning solutions of the twistor equation (see [BFGK]). For example, in [Li] (see also proposition 6) one can find the following theorem: if $f : M^2 \to \mathbb{R}^1$ is a real-valued function such that the equation

$$\nabla_{\vec{T}}(\varphi) + \frac{1}{2} f \cdot \vec{T} \cdot \varphi = 0$$

admits a non-trivial solution then f is constant and $f^2 = G$. In the theory of surfaces this statement correspondends to the fact that an umbilic surface is a part of the sphere or the plane. Indeed, an umbilic surface $M^2 \hookrightarrow \mathbb{R}^3$ admits a spinor field φ such that

$$\nabla_{\vec{T}}(\varphi) + \frac{1}{2} H \vec{T} \cdot \varphi = 0$$

and therefore $H^2 = G = const$, i.e. the second fundamental form is proportional to the metric. In a similar way one can translate other facts of the theory of surfaces into properties of solutions of the equation $\nabla_X \varphi = E(X) \cdot \varphi$.

References

[Bä] C. Bär. Extrinsic bounds for eigenvalues of the Dirac operator, Preprint Universität Freiburg, Oktober 1997.

[BFGK] H. Baum, Th. Friedrich, R. Grunewald, I. Kath. Twistors and Killing spinors on Riemannian manifolds, Teubner-Verlag Leipzig / Stuttgart 1991.

[F] Th. Friedrich. Dirac-Operatoren in der Riemannschen Geometrie, Vieweg-Verlag Braunschweig/ Wiesbaden 1997.

[KPP] G. Kamberov, F. Pedit, U. Pinkall. Bonnet pairs and isothermic surfaces, Preprint dg-ga 9610006.

[KS] R. Kussner, N. Schmitt. Representation of surfaces in space, Preprint dg-ga 9610005.

[La] H.B. Lawson. The global behaviour of minimal surfaces in S^n, Ann. Math. 1970, V. 92, No.2, 224-237.

[Li] A. Lichnerowicz. Spin manifolds, Killing spinors and universality of the Hijazi inequality, Lett. Math. Phys. 13 (1987), 331–344.

[R] J. Richter. Conformal maps of a Riemannian surface into the space of quaternions, Dissertation TU Berlin 1997.

[S] A. Svec. Global differential geometry of surfaces, Verlag der Wissenschaften, Berlin 1981.

[Tai1] I.A. Taimanov. Surfaces of revolution in terms of solutions, Ann. Glob. Anal. Geom. 15 (1997), 419-435.

[Tai2] I.A. Taimanov. Modified Novikov-Veselov equation and differential geometry of surfaces, Transl. Amer. Math. Soc., Ser. 2, V. 179 (1997), 133-151.

[Tai3] I.A. Taimanov. The Weierstraß representation of closed surfaces in $\mathbb{R}^3$, Preprint SFB 288, No. 291, TU Berlin (1997), dg-ga 9710029.

[Tr1] A. Trautman. Spinors and the Dirac operator on hypersurfaces I. General Theory, Journ. Math. Phys. 33 (1992), 4011–4019.

[Tr2] A. Trautman. The Dirac operator on hypersurfaces, Acta Phys. Polon. B 26 (1995), 1283–1310.

Thomas Friedrich
Humboldt-Universität zu Berlin
Institut für Reine Mathematik
Sitz: Ziegelstraße 13a
D-10099 Berlin
e-mail: friedric@mathematik.hu-berlin.de